arXiv:1605.02484v1 [math.FA] 9 May 2016

ON REFINED OPERATOR VERSION OF YOUNG INEQUALITY AND ITS REVERSE

ALEMEH SHEIKHHOSSEINI[1] MARYAM KHOSRAVI[2*]

Abstract. In this note, some refinements of Young inequality and its reverse for positive numbers are proved and using these inequalities some operator versions and Hilbert-Schmidt norm versions for matrices of these inequalities are obtained.

1. Introduction

Let $\mathbb{B}(\mathcal{H})$ denote the C^*-algebra of all bounded linear operators on a complex Hilbert space $\mathcal{H}$. In the case when dim $\mathcal{H} = n$, we identify $\mathbb{B}(\mathcal{H})$ with the matrix algebra $\mathbb{M}_n$ of all $n \times n$ complex matrices. For $A = (a_{ij}) \in \mathbb{M}_n$, the Hilbert-Schmidt norm of A is defined by

$$\|A\|_2 = \left(\sum_{i,j=1}^{n} |a_{ij}|^2\right)^{\frac{1}{2}} = \sum_{i=1}^{n} \left(s_j^2(A)\right)^{\frac{1}{2}},$$

where $s_j(A)$ $(1 \leq j \leq n)$ are the singular values of A. It is known that $\|.\|_2$ is a unitarily invariant norm.

Let $A, B \in \mathbb{M}_n$. Denoted by $A \circ B$ the Schur (Hadamard) product of A and B, that is, the entrywise product.

For positive real numbers a and b, the classical Young inequality says that if $\nu \in [0, 1]$, then

$$a^{1-\nu} b^{\nu} \leqslant (1 - \nu)a + \nu b,$$

with equality if and only if $a = b$. When $\nu = \frac{1}{2}$, the Young inequality is called the arithmetic-geometric mean inequality

$$\sqrt{ab} \leqslant \frac{a + b}{2}. \tag{1.1}$$

2010 *Mathematics Subject Classification.* Primary: 47A63; Secondary: 47A64, 15A42.

Key words and phrases. Young inequality, positive operators, weighted means, Hilbert-Schmidt norm

∗ Corresponding author.

Throughout, we denote $a^{1-\nu}b^{\nu}$ and $(1-\nu)a+\nu b$, respectively by $a\sharp_{\nu}b$ and $a\nabla_{\nu}b$. The Heinz mean is defined as

$$H_{\nu}(a,b)=\frac{a^{1-\nu}b^{\nu}+a^{\nu}b^{1-\nu}}{2}$$

for $a,b\geqslant 0$ and $\nu\in[0,1]$. It's easy to see that

$$\sqrt{ab}\leqslant H_{\nu}(a,b)\leqslant\frac{a+b}{2}.$$

In [9] and [10], F. Kittaneh and Y. Manasrah improved the Young inequality and its reverse as follows:

$$a^{1-\nu}b^{\nu}+r(\sqrt{a}-\sqrt{b})^2\leqslant(1-\nu)a+\nu b\leqslant a^{1-\nu}b^{\nu}+s(\sqrt{a}-\sqrt{b})^2, \tag{1.2}$$

where $r=\min\{\nu,1-\nu\}$ and $s=\max\{\nu,1-\nu\}$.

The authors of [7] and [8] obtained another refinement of the Young inequality as follows:

$$r^2(a-b)^2\leqslant((1-\nu)a+\nu b)^2-(a^{1-\nu}b^{\nu})^2\leqslant s^2(a-b)^2, \tag{1.3}$$

where $r=\min\{\nu,1-\nu\}$, and $s=\max\{\nu,1-\nu\}$.

Recently, J. Zhao and J. Wu [13] obtained the following refinement of inequality (1.2):

$$\begin{aligned}r((ab)^{\frac{1}{4}}-\sqrt{a})^2+\nu(\sqrt{a}-\sqrt{b})^2&\leqslant(1-\nu)a+\nu b-a^{1-\nu}b^{\nu}\\&\leqslant(1-\nu)(\sqrt{a}-\sqrt{b})^2-r((ab)^{\frac{1}{4}}-\sqrt{b})^2,\end{aligned}$$

where $0\leqslant\nu\leqslant\frac{1}{2}$ and $r=\min\{2\nu,1-2\nu\}$ and

$$\begin{aligned}r((ab)^{\frac{1}{4}}-\sqrt{b})^2+(1-\nu)(\sqrt{a}-\sqrt{b})^2&\leqslant(1-\nu)a+\nu b-a^{1-\nu}b^{\nu}\\&\leqslant\nu(\sqrt{a}-\sqrt{b})^2-r((ab)^{\frac{1}{4}}-\sqrt{a})^2,\end{aligned}$$

where $\frac{1}{2}\leqslant\nu\leqslant 1$ and $r=\min\{2(1-\nu),1-2(1-\nu)\}$.
Also, they obtained the following refinement of inequalities (1.3):

$$\begin{aligned}r(\sqrt{ab}-a)^2+\nu^2(a-b)^2&\leqslant((1-\nu)a+\nu b)^2-(a^{1-\nu}b^{\nu})^2\\&\leqslant(1-\nu)^2(a-b)^2-r(\sqrt{ab}-b)^2,\end{aligned} \tag{1.4}$$

where $0\leqslant\nu\leqslant\frac{1}{2}$ and $r=\min\{2\nu,1-2\nu\}$ and

$$\begin{aligned}r(\sqrt{ab}-b)^2+(1-\nu)^2(a-b)^2&\leqslant((1-\nu)a+\nu b)^2-(a^{1-\nu}b^{\nu})^2\\&\leqslant\nu^2(a-b)^2-r(\sqrt{ab}-a)^2,\end{aligned} \tag{1.5}$$

where $\frac{1}{2}\leqslant\nu\leqslant 1$ and $r=\min\{2(1-\nu),1-2(1-\nu)\}$.

Let $A,B\in\mathbb{B}(\mathcal{H})$ be two operators and $\nu\in[0,1]$. The $\nu-$weighted arithmetic mean of A and B, denoted and defined by:

$$A\nabla_{\nu}B=(1-\nu)A+\nu B.$$

If A is invertible, $\nu-$geometric mean and $\nu-$Heinz mean of A and B are defined respectively, as

$$A\sharp_\nu B = A^{\frac{1}{2}}(A^{-\frac{1}{2}}BA^{-\frac{1}{2}})^\nu A^{\frac{1}{2}}$$

and

$$H_\nu(A,B) = \frac{A\sharp_\nu B + A\sharp_{1-\nu} B}{2}.$$

In addition, if both A and B are invertible, $\nu-$harmonic mean of A and B, denoted by $A!_\nu B$, is defined as

$$A!_\nu B = ((1-\nu)A^{-1} + \nu B^{-1})^{-1}.$$

When $\nu = \frac{1}{2}$, we write $A\nabla B, A\sharp B$ and $A!B$ for brevity, respectively.
It is well known that if A and B are positive invertible operators, then

$$A\nabla_\nu B \geqslant A\sharp_\nu B \geqslant A!_\nu B,$$

for $0 < \nu < 1$; see [4, 6] for more information.

Based on the refined Young inequality (1.4) and its reverse (1.5), J. Zhao and J. Wu [13] proved that if $A, B, X \in \mathbb{M}_n$ such that A and B are positive semidefinite, then

$$\begin{aligned}\nu^2\|AX - XB\|_2^2 + r\|A^{\frac{1}{2}}XB^{\frac{1}{2}} - AX\|_2^2 & \\ \leqslant \|(1-\nu)AX + \nu XB\|_2^2 - \|A^{1-\nu}XB^\nu\|_2^2 & \\ \leqslant (1-\nu)^2\|AX - XB\|_2^2 + -r\|A^{\frac{1}{2}}XB^{\frac{1}{2}} - XB\|_2^2, & \quad (1.6)\end{aligned}$$

where $0 \leqslant \nu \leqslant \frac{1}{2}$ and $r = \min\{2\nu, 1-2\nu\}$ and

$$\begin{aligned}(1-\nu)^2\|AX - XB\|_2^2 + r\|A^{\frac{1}{2}}XB^{\frac{1}{2}} - XB\|_2^2 & \\ \leqslant \|(1-\nu)AX + \nu XB\|_2^2 - \|A^{1-\nu}XB^\nu\|_2^2 & \\ \leqslant \nu^2\|AX - XB\|_2^2 - r\|A^{\frac{1}{2}}XB^{\frac{1}{2}} - AX\|_2^2, & \quad (1.7)\end{aligned}$$

where $\frac{1}{2} \leqslant \nu \leqslant 1$ and $r = \min\{2(1-\nu), 1-2(1-\nu)\}$.

Their results were generalized by Liao and Wu [11], using Kantorovich constant.

Furthermore, some similar results can be found in [1, 3].

In addition, in [2], the authors investigated on these inequalities, for the cases that $\nu \leq 0$ or $\nu \geq 1$. In these cases, they proved the reverse of some of these inequalities. Furthermore, in [12], the numerical version of some of these relations, are discussed.

The main aim of this paper, is to state a generalization of these inequalities. First, we present some generalizations of numerical inequalities and base of them we prove some refined operator versions of Young inequality and its reverse. Also some inequalities for Hilbert-Schmidt norm of matrices are obtained.

In this paper, for $0 < \nu < 1$, the notations $m_k = \lfloor 2^k\nu \rfloor$ is the largest integer not greater than $2^k\nu$, $r_0 = \min\{\nu, 1-\nu\}$ and $r_k = \min\{2r_{k-1}, 1 - 2r_{k-1}\}$, for $k \geq 1$.

2. Numerical results

We start with some numerical results.

Theorem 2.1. *Let a, b be two positive real numbers and $\nu \in (0,1)$. Then*

$$a\nabla_\nu b \geqslant a\sharp_\nu b + \sum_{k=0}^{\infty} r_k \big[\big(a^{1-\frac{m_k}{2^k}} b^{\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(a^{1-\frac{m_k+1}{2^k}} b^{\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}}\big]^2. \tag{2.1}$$

In addition, if $\nu = \frac{t}{2^n}$ for some $t, n \in \mathbb{N}$, then

$$a\nabla_\nu b = a\sharp_\nu b + \sum_{k=0}^{n-1} r_k \big[\big(a^{1-\frac{m_k}{2^k}} b^{\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(a^{1-\frac{m_k+1}{2^k}} b^{\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}}\big]^2.$$

Proof. It is enough to prove that for each $n \in \mathbb{N} \cup \{0\}$,

$$a\nabla_\nu b \geqslant a\sharp_\nu b + \sum_{k=0}^{n} r_k \big[\big(a^{1-\frac{m_k}{2^k}} b^{\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(a^{1-\frac{m_k+1}{2^k}} b^{\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}}\big]^2. \tag{2.2}$$

We prove it by induction. For $n = 0$, we get to the well-known inequality (1.2). Let inequality (2.2) holds for n.

First, let $0 < \nu < \frac{1}{2}$. Thus, we have

$$\begin{aligned} a\nabla_\nu b - r_0(\sqrt{a} - \sqrt{b})^2 &= a\nabla_\nu b - \nu(\sqrt{a} - \sqrt{b})^2 \\ &= 2\nu\sqrt{ab} + (1 - 2\nu)a \\ &= a\nabla_{2\nu}\sqrt{ab} \end{aligned}$$

Applying inequality (2.2) for two positive numbers a and $\sqrt{ab}$ and $2\nu \in (0,1)$, we have

$$\begin{aligned} a\nabla_\nu b - r_0(\sqrt{a} - \sqrt{b})^2 &= a\nabla_{2\nu}\sqrt{ab} \\ &\geq a\sharp_{2\nu}\sqrt{ab} + \sum_{k=0}^{n} r_{k+1}\big[\big(a^{1-\frac{m_{k+1}}{2^k}} (\sqrt{ab})^{\frac{m_{k+1}}{2^k}}\big)^{\frac{1}{2}} \\ &\quad - \big(a^{1-\frac{m_{k+1}+1}{2^k}} (\sqrt{ab})^{\frac{m_{k+1}+1}{2^k}}\big)^{\frac{1}{2}}\big]^2 \\ &= a\sharp_\nu b + \sum_{k=1}^{n+1} r_k \big[\big(a^{1-\frac{m_k}{2^k}} b^{\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(a^{1-\frac{m_k+1}{2^k}} b^{\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}}\big]^2. \end{aligned}$$

For $\frac{1}{2} < \nu < 1$, we can apply the first part for $1 - \nu$ and replace a and b. Note that $[2^k(1-\nu)] = 2^k - [2^k\nu] - 1$ if $2^k\nu$ is not integer. Thus, if $2^k\nu$ is not integer for each k, the inequality follows.

Now, let $\nu = \frac{t}{2^\ell}$ for some odd number t and $\ell \in \mathbb{N} \cup \{0\}$. Since for each $i < \ell$, the coefficient $r_i \leq \frac{1}{2}$ is of the form $\frac{t_i}{2^{\ell-i}}$, it can be concluded that $r_\ell = 0$ and so $r_k = 0$ for all $k \geq n$. On the other hand $2^k\nu$ is not integer for $k < \ell$. So the result follows.

A similar argument, shows the equality holds when $\nu = \frac{t}{2^n}$. □

Remark 2.2. Note that the series appear in this theorem is a positive series with a finite upper bound. So it is convergent. This fact is also satisfies with all other series appear in this note.

Changing the place of numbers a and b in inequality (2.1), we can state the following result for Heinz mean.

Corollary 2.3. *Let a, b be two positive real numbers and $\nu \in (0,1)$. Then*

$$a\nabla b \geqslant H_\nu(a,b) + \sum_{k=0}^{\infty} r_k \big[H_{\frac{m_k}{2^k}}(a,b) - 2H_{\frac{2m_k+1}{2^{k+1}}}(a,b) + H_{\frac{m_k+1}{2^k}}(a,b)\big].$$

In the following theorem, we state a reverse of Young inequality.

Theorem 2.4. *Let a, b be two positive real numbers and $\nu \in (0,1)$. Then*

$$a\nabla_\nu b \leqslant a\sharp_\nu b + (\sqrt{a}-\sqrt{b})^2 - \sum_{k=0}^{\infty} r_k \big[\big(a^{\frac{m_k}{2^k}} b^{1-\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(a^{\frac{m_k+1}{2^k}} b^{1-\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}}\big]^2. \tag{2.3}$$

Proof. By $a\sharp_\nu b + b\sharp_\nu a \geq 2\sqrt{ab}$, and inequality (2.1), we have

$$\begin{aligned}(\sqrt{a}-\sqrt{b})^2 - a\nabla_\nu b &= b\nabla_\nu a - 2\sqrt{ab} \\ &\geq -a\sharp_\nu b + \sum_{k=0}^{\infty} r_k \big[\big(a^{\frac{m_k}{2^k}} b^{1-\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(a^{\frac{m_k+1}{2^k}} b^{1-\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}}\big]^2.\end{aligned}$$

So the result follows. □

Corollary 2.5. *Let a, b be two positive real numbers and $\nu \in (0,1)$. Then*

$$a\nabla b \leqslant H_\nu(a,b) + (\sqrt{a}-\sqrt{b})^2 - \sum_{k=0}^{\infty} r_k \big[H_{\frac{m_k}{2^k}}(a,b) - 2H_{\frac{2m_k+1}{2^{k+1}}}(a,b) + H_{\frac{m_k+1}{2^k}}(a,b)\big].$$

Remark 2.6. Replacing a and b by their squares in (2.1) and (2.3), respectively, we obtain

$$a^2\nabla_\nu b^2 \geqslant a^2\sharp_\nu b^2 + \sum_{k=0}^{\infty} r_k \big[a^{1-\frac{m_k}{2^k}} b^{\frac{m_k}{2^k}} - a^{1-\frac{m_k+1}{2^k}} b^{\frac{m_k+1}{2^k}}\big]^2 \tag{2.4}$$

and

$$a^2\nabla_\nu b^2 \leqslant a^2\sharp_\nu b^2 + (a-b)^2 - \sum_{k=0}^{\infty} r_k \big[a^{\frac{m_k}{2^k}} b^{1-\frac{m_k}{2^k}} - a^{\frac{m_k+1}{2^k}} b^{1-\frac{m_k+1}{2^k}}\big]^2. \tag{2.5}$$

The following two theorems, are useful to prove a version of these inequalities for the Hilbert-Schmidt norm of matrices.

Theorem 2.7. *Let a, b be two positive real numbers and $\nu \in (0,1)$. Then*

$$(a\nabla_\nu b)^2 \geqslant (a\sharp_\nu b)^2 + r_0^2(a-b)^2 + \sum_{k=1}^{\infty} r_k \big[a^{1-\frac{m_k}{2^k}} b^{\frac{m_k}{2^k}} - a^{1-\frac{m_k+1}{2^k}} b^{\frac{m_k+1}{2^k}}\big]^2. \tag{2.6}$$

Proof. By (2.4), we have

$$(a\nabla_\nu b)^2 - r_0^2(a-b)^2 = a^2\nabla_\nu b^2 - r_0(a-b)^2$$
$$\geq (a\sharp_\nu b)^2 + \sum_{k=1}^{\infty} r_k\big[a^{1-\frac{m_k}{2^k}}b^{\frac{m_k}{2^k}} - a^{1-\frac{m_k+1}{2^k}}b^{\frac{m_k+1}{2^k}}\big]^2$$

□

Theorem 2.8. *Let a, b be two positive real numbers and $\nu \in (0,1)$. Then*

$$(a\nabla_\nu b)^2 \leqslant (a\sharp_\nu b)^2 + (1-r_0)^2(a-b)^2 - \sum_{k=1}^{\infty} r_k\big[a^{1-\frac{m_k}{2^k}}b^{\frac{m_k}{2^k}} - a^{1-\frac{m_k+1}{2^k}}b^{\frac{m_k+1}{2^k}}\big]^2. \tag{2.7}$$

Proof. We have

$$\begin{aligned}(a\nabla_\nu b)^2 &- (1-r_0)^2(a-b)^2\\ &= a^2\nabla_\nu b^2 - (1-r_0)(a-b)^2\\ &\leqslant (a\sharp_\nu b)^2 + r_0(a-b)^2\\ &\quad - \sum_{k=0}^{\infty} r_k\big[a^{1-\frac{m_k}{2^k}}b^{\frac{m_k}{2^k}} - a^{1-\frac{m_k+1}{2^k}}b^{\frac{m_k+1}{2^k}}\big]^2\\ &\qquad\qquad \text{by inequality (2.5)}\\ &= (a\sharp_\nu b)^2 - \sum_{k=1}^{\infty} r_k\big[a^{1-\frac{m_k}{2^k}}b^{\frac{m_k}{2^k}} - a^{1-\frac{m_k+1}{2^k}}b^{\frac{m_k+1}{2^k}}\big]^2.\end{aligned}$$

□

3. Related operator inequalities

Two state the operator versions of the inequalities obtained in section 2, we need the following lemma.

Lemma 3.1. [5] *Let $X \in \mathbb{B}(\mathcal{H})$ be self-adjoint and let f and g be continuous real functions such that $f(t) \geqslant g(t)$ for all $t \in \sigma(X)$ (the spectrum of X). Then $f(X) \geqslant g(X)$.*

Next, we give the first result in this section, which is based on Theorem 2.1 and is a refinement of Theorem 1 in [13].

Theorem 3.2. *Let $A, B \in \mathbb{B}(\mathcal{H})$ be two positive invertible operators and $\nu \in (0,1)$.*

$$A\nabla_\nu B \geqslant A\sharp_\nu B + \sum_{k=0}^{\infty} r_k[A\sharp_{\frac{m_k}{2^k}} B - 2A\sharp_{\frac{2m_k+1}{2^{k+1}}} B + A\sharp_{\frac{m_k+1}{2^k}} B]. \tag{3.1}$$

Proof. Choosing $a = 1$, in Theorem 2.1, we have

$$1 - \nu + \nu b \geqslant b^{\nu} + \sum_{k=0}^{\infty} r_k \big[\big(b^{\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(b^{\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}} \big]^2,$$

for any $b > 0$.
If $X = A^{-\frac{1}{2}} B A^{-\frac{1}{2}}$, then $\sigma(X) \subseteq (0, \infty)$. According to Lemma 3.1, we get

$$(1-\nu)I + \nu X \geqslant X^{\nu} + \sum_{k=0}^{\infty} r_k [X^{\frac{m_k}{2^k}} - 2X^{\frac{2m_k+1}{2^{k+1}}} + X^{\frac{m_k+1}{2^k}}].$$

Multiplying both sides by $A^{\frac{1}{2}}$, we obtain

$$A \nabla_{\nu} B \geqslant A \sharp_{\nu} B + \sum_{k=0}^{\infty} r_k [A \sharp_{\frac{m_k}{2^k}} B - 2A \sharp_{\frac{2m_k+1}{2^{k+1}}} B + A \sharp_{\frac{m_k+1}{2^k}} B].$$

This completes the proof. □

Since for all positive integer n,

$$f(t) = \sum_{k=0}^{n} r_k \big[\big(t^{\frac{m_k}{2^k}}\big)^{\frac{1}{2}} - \big(t^{\frac{m_k+1}{2^k}}\big)^{\frac{1}{2}} \big]^2 = \sum_{k=0}^{n} r_k \big[t^{\frac{m_k}{2^k}} - 2t^{\frac{2m_k+1}{2^{k+1}}} + t^{\frac{m_k+1}{2^k}} \big]$$

is a continuous function on $[0, \infty)$ and $A^{-\frac{1}{2}} B A^{-\frac{1}{2}}$ is a positive operator, then $\sigma(f(A^{-\frac{1}{2}} B A^{-\frac{1}{2}})) \subseteq [0, \infty)$. Thus

$$A^{\frac{1}{2}} f(A^{-\frac{1}{2}} B A^{-\frac{1}{2}}) A^{\frac{1}{2}} = \sum_{k=0}^{n} r_k (A \sharp_{\frac{m_k}{2^k}} B - 2A \sharp_{\frac{2m_k+1}{2^{k+1}}} B + A \sharp_{\frac{m_k+1}{2^k}} B),$$

is a positive operator. Then by inequality (3.1), we obtain

$$A \sharp_{\nu} B \leqslant A \sharp_{\nu} B + \sum_{k=0}^{n} r_k (A \sharp_{\frac{m_k}{2^k}} B - 2A \sharp_{\frac{2m_k+1}{2^{k+1}}} B + A \sharp_{\frac{m_k+1}{2^k}} B) \leqslant A \nabla_{\nu} B.$$

and therefore

$$A \sharp_{\nu} B \leqslant A \sharp_{\nu} B + \sum_{k=0}^{\infty} r_k (A \sharp_{\frac{m_k}{2^k}} B - 2A \sharp_{\frac{2m_k+1}{2^{k+1}}} B + A \sharp_{\frac{m_k+1}{2^k}} B) \leqslant A \nabla_{\nu} B. \tag{3.2}$$

Replacing A and B by A^{-1} and B^{-1} respectively, we obtain

$$\begin{aligned} & A^{-1} \sharp_{\nu} B^{-1} \\ & \leqslant A^{-1} \sharp_{\nu} B^{-1} + \sum_{k=0}^{\infty} r_k (A^{-1} \sharp_{\frac{m_k}{2^k}} B^{-1} - 2A^{-1} \sharp_{\frac{2m_k+1}{2^{k+1}}} B^{-1} + A^{-1} \sharp_{\frac{m_k+1}{2^k}} B^{-1}) \\ & \leqslant A^{-1} \nabla_{\nu} B^{-1}. \end{aligned} \tag{3.3}$$

Taking inverse in (3.3), we have

$$\begin{aligned} &A!_\nu B \\ &\leqslant \{A^{-1}\sharp_\nu B^{-1} + \sum_{k=0}^{n} r_k(A^{-1}\sharp_{\frac{m_k}{2^k}} B^{-1} - 2A^{-1}\sharp_{\frac{2m_k+1}{2^{k+1}}} B^{-1} + A^{-1}\sharp_{\frac{m_k+1}{2^k}} B^{-1})\}^{-1} \\ &\leqslant A\sharp_\nu B. \end{aligned} \tag{3.4}$$

It is worth to mention that inequalities (3.2), (3.3) and (3.4) are respectively refinements of inequalities (30)-(34) in [13].
The following theorem is an operator version of Theorem 2.4 and is a refinement of Theorem 2 in [13].

Theorem 3.3. *Let $A, B \in \mathbb{B}(\mathcal{H})$ be two positive invertible operators and $\nu \in (0,1)$.*

$$A\nabla_\nu B \leqslant A\sharp_\nu B + (A - 2A\sharp B + B) - \sum_{k=0}^{\infty} r_k[A\sharp_{\frac{m_k}{2^k}} B - 2A\sharp_{\frac{2m_k+1}{2^{k+1}}} B + A\sharp_{\frac{m_k+1}{2^k}} B].$$

Proof. By Theorem 2.4, using the same ideas as in the proof of Theorem 3.2, we can get the result. □

Corollary 3.4. *Let $A, B \in \mathbb{B}(\mathcal{H})$ be two positive invertible operators and $\nu \in (0,1)$. Then*

$$A\nabla B \geqslant H_\nu(A,B) + \sum_{k=0}^{\infty} r_k[H_{\frac{m_k}{2^k}}(A,B) - 2H_{\frac{2m_k+1}{2^{k+1}}}(A,B) + H_{\frac{m_k+1}{2^k}}(A,B)].$$

and

$$\begin{aligned} A\nabla B \leqslant H_\nu(A,B) &+ (A - 2A\sharp B + B) \\ &- \sum_{k=0}^{\infty} r_k[H_{\frac{m_k}{2^k}}(A,B) - 2H_{\frac{2m_k+1}{2^{k+1}}}(A,B) + H_{\frac{m_k+1}{2^k}}(A,B)]. \end{aligned}$$

4. Hilbert-Schmidt norm version

In this section, we obtain some inequalities for the Hilbert-Schmidt norm. Applying Theorem 2.7, we get the following theorem that is a refinement of first inequalities in (1.6) and (1.7).

Theorem 4.1. *Let $A, B, X \in \mathbb{M}_n$ such that A and B are two positive semi-definite matrices and $\nu \in (0,1)$. Then*

$$\begin{aligned} \|A^{1-\nu}XB^\nu\|_2^2 + r_0^2\|AX - XB\|_2^2 + \sum_{k=1}^{\infty} r_k \|A^{1-\frac{m_k}{2^k}} X B^{\frac{m_k}{2^k}} - A^{1-\frac{m_k+1}{2^k}} X B^{\frac{m_k+1}{2^k}}\|_2^2 \\ \leqslant \|(1-\nu)AX - \nu XB\|_2^2. \end{aligned}$$

Proof. Since A and B are positive semidefinite $n \times n$ matrices, there exist unitary matrices $U, V \in \mathbb{M}_n$ such that $A = Udiag(\lambda_1, \ldots, \lambda_n)U^*$ and $B = Vdiag(\mu_1, \ldots, \mu_n)V^*$. Let $Y = U^*XV = (y_{ij})$. Then it's straightforward to check that

$$(1-\nu)AX - \nu XB = U[((1-\nu)\lambda_i + \nu\mu_j) \circ Y]V^*,$$

$$AX - XB = U[(\lambda_i - \mu_j) \circ Y]V^*$$

$$A^{1-\nu}XB^{\nu} = U[(\lambda_i^{1-\nu}\mu_j^{\nu}) \circ Y]V^*$$

and

$$A^{1-\frac{m_k}{2^k}}XB^{\frac{m_k}{2^k}} - A^{1-\frac{m_k+1}{2^k}}XB^{\frac{m_k+1}{2^k}} = U[(\lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}} - \lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}}) \circ Y]V^*.$$

Utilizing the unitarily invariant property of $\|.\|_2$ and Theorem 2.7, we have

$$\begin{aligned}
&\|(1-\nu)AX - \nu XB\|_2^2 \\
&= \|((1-\nu)\lambda_i + \nu\mu_j) \circ Y\|_2^2 \\
&= \sum_{i,j=1}^{n} ((1-\nu)\lambda_i + \nu\mu_j)^2 |y_{ij}|^2 \\
&\geqslant \sum_{i,j=1}^{n} \left\{ (\lambda_i^{1-\nu}\mu_j^{\nu})^2 + r_0^2(\lambda_i - \mu_j)^2 + \sum_{k=1}^{\infty} r_k(\lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}} - \lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}})^2 \right\} |y_{ij}|^2 \\
&= \sum_{i,j=1}^{n} (\lambda_i^{1-\nu}\mu_j^{\nu})^2 |y_{ij}|^2 + \sum_{i,j=1}^{n} r_0^2(\lambda_i - \mu_j)^2 |y_{ij}|^2 \\
&\quad + \sum_{i,j=1}^{n} \left\{ \sum_{k=1}^{\infty} r_k(\lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}} - \lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}})^2 |y_{ij}|^2 \right\} \\
&= \sum_{i,j=1}^{n} (\lambda_i^{1-\nu}\mu_j^{\nu})^2 |y_{ij}|^2 + \sum_{i,j=1}^{n} r_0^2(\lambda_i - \mu_j)^2 |y_{ij}|^2 \\
&\quad + \sum_{k=1}^{\infty} \left\{ \sum_{i,j=1}^{n} r_k(\lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}} - \lambda_i^{1-\frac{m_k}{2^k}}\mu_j^{\frac{m_k}{2^k}})^2 |y_{ij}|^2 \right\} \\
&= \|A^{1-\nu}XB^{\nu}\|_2^2 + r_0^2\|AX - XB\|_2^2 \\
&\quad + \sum_{k=1}^{\infty} r_k \|A^{1-\frac{m_k}{2^k}}XB^{\frac{m_k}{2^k}} - A^{1-\frac{m_k+1}{2^k}}XB^{\frac{m_k+1}{2^k}}\|_2^2.
\end{aligned}$$

So, the proof is complete. □

The last theorem is a refinement of second inequalities in (1.6) and (1.7).

Theorem 4.2. *Let $A, B, X \in \mathbb{M}_n$ such that A and B are two positive semi-definite matrices and $\nu \in (0,1)$. Then*

$$\|(1-\nu)AX - \nu XB\|_2^2 \leqslant \|A^{1-\nu}XB^{\nu}\|_2^2 + (1-r_0)^2\|AX - XB\|_2^2 - \sum_{k=1}^{\infty} r_k \|A^{1-\frac{m_k}{2^k}} XB^{\frac{m_k}{2^k}} - A^{1-\frac{m_k+1}{2^k}} XB^{\frac{m_k+1}{2^k}}\|_2^2.$$

Proof. By Theorem 2.8 and using the same idea as in the proof of Theorem 4.1, we can obtain the desired result. □

References

1. H. Alzer, C. M. da Fonseca and A. Kovačec, *Young-type inequalities and their matrix analogues*, Linear Multilinear Algebra. **63** (2015), no. 3, 622–635.
2. M. Bakherad, M. Krnic and M.S. Moslehian, *Reverses of the Young inequality for matrices and operators*, Rocky Mountain J. Math. (to appear).
3. Y. Feng, *Refinements of Young inequalities for matrices*, Far. East. J. Math. Sci. **63** (2012), 45–55.
4. T. Furuta, *Invitation to Linear Operators: Form Matrix to Bounded Linear Operators on a Hilbert Space*, Taylor and Francis, 2002.
5. T. Furuta, J. Mićić Hot and J. Pečarić, *Mond–Pečarić Method in Operator Inequalities*, Element, Zagreb, 2005.
6. T. Furuta and M. Yanagide, *Generalized means and convexity of inversion for positive operators*, Amer. Math. Monthly 105 (1998) 258-259.
7. C.J. He and L.M. Zou, *Some inequalities involving unitarily invariant norms*, Math. Inequal. Appl. 12 (4) (2012) 767-776.
8. O. Hirzallah and F. Kittaneh, *Matrix Young inequalities for the Hilbert-Schmidt norm*, Linear Algebra Appl. 308 (2000) 77-84.
9. F. Kittaneh and Y. Manasrah, *Improved Young and Heinz inequalities for matrices*, J. Math. Anal. Appl. 361 (2010) 262-269.
10. F. Kittaneh and Y. Manasrah, *Reverse Young and Heinz inequalities for matrices*, Linear Multilinear Algebra 59 (9) (2011) 1031-1037.
11. W. Liao and J. Wu, *Improved Young and Heinz inequalities with the Kantorovich constant,* , J. Math. Inequal., Preprint.
12. A. Salemi and A. Sheikh Hosseini, *Matrix Young numerical radius inequalities*, Math. Inequal. Appl. 16 (2013), no. 3, 783-791.
13. J. Zhao and J. Wu, *Operator inequalities involving improved Young and its reverse inequalities*, J. Math. Anal. Appl. 421 (2015) 1779-1789.

[1] Department of Pure Mathematics, Faculty of Mathematics and Computer, Shahid Bahonar University of Kerman, Kerman, Iran
E-mail address: sheikhhosseini@uk.ac.ir; hosseini8560@gmail.com

[2] Department of Pure Mathematics, Faculty of Mathematics and Computer, Shahid Bahonar University of Kerman, Kerman, Iran
E-mail address: khosravi_m @uk.ac.ir; khosravi_m2000@yahoo.com